



Genemed Synthesis, Inc.  
6203 Wood Lake Center Dr., Bldg. 2, San Antonio, TX 78244, USA  
**Toll free** (800) 344-5337; **Phone:** (210) 745-5988; **Fax** (210) 745-5992  
**Email:** [info@genemedsyn.com](mailto:info@genemedsyn.com), **Website:** [www.genemedsyn.com](http://www.genemedsyn.com)

## Raji Cell Lysate

**Catalog No:** CL-66-1207

**Size:** 100 µg

**Cell line:** Raji (Human Burkitt's lymphoma)

**Source:** Human (11yrs, Male) B lymphocyte Burkitt's lymphoma

**Growth media:** RPMI & 10% NCS (Newborn calf serum)

### Cell lysate preparation:

Raji cell lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The cell lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% β-mercaptoethanol.

### Application:

Raji cell lysate is ready to load on SDS-PAGE for Western blotting, 20 µg per lane is recommended for mini gel.

### Storage:

It is supplied as whole cell lysate at 100 µg at 2 mg/ml in 1 x SDS sample buffer containing 5% β-mercaptoethanol. Store at -20°C for three months or at -70°C for one year.

### Usage:

This item is for LABORATORY RESEARCH USE ONLY.

Rev. 120822V